arXiv:0904.0906v1 [math.RA] 6 Apr 2009

A short proof of Grinshpon's theorem

Dinesh Khurana

Indian Institute of Science Education and Research, Mohali
MGSIPA Transit Campus, Sector 26 Chandigarh-160019, India
`dkhurana@iisermohali.ac.in`

In [G, Theorem], Grinshpon proved that if S is a commutative subring of a ring R and $A \in M_n(S)$ is invertible in $M_n(R)$, then $det(A)$ is invertible in R. Grinshpon's theorem is immediate from the following result by taking $B = A^{-1}$ and using the fact that a square matrix over a commutative ring is invertible iff its determinant is invertible. For $X \in M_n(R)$, let $C(X) = \{Y \in M_n(R) : XY = YX\}$.

Theorem. *Let A, $B \in M_n(R)$ be such that $C(A) = C(B)$. If the entries of A commute with each other, then there exists a commutative subring T of R such that $A, B \in M_n(T)$.*

Proof. Let I denote the identity of $M_n(R)$. If a is any entry of A, then $aI \in C(A) = C(B)$. Thus every entry of A commutes with every entry of B. So if b is any entry of B, then $bI \in C(A) = C(B)$ implying that the entries of B commute with each other also. Thus the subring T generated by all entries of A and B is commutative. $\square$

References

[G] M. Grinshpon, *Invertibility of matrices over subrings*, communications in Algebra 36 (2008), 2619-2624.